

NOTES ON GEOGRAPHIC DISTRIBUTION

Amphibia, Centrolenidae, *Cochranella pulverata*, *Hyalinobatrachium aureoguttatum*:
Distribution extension, Ecuador.

Martín R. Bustamante ¹
Diego F. Cisneros-Heredia ^{2, 3, 4}
Mario H. Yáñez-Muñoz ⁴
H. Mauricio Ortega-Andrade ⁴
Juan M. Guayasamin ^{1, 5}

¹ Museo de Zoología, Centro de Biodiversidad y Ambiente, Escuela de Biología, Pontificia Universidad Católica del Ecuador, Apartado 17-01-2184, Quito, Ecuador. E-mail: mrbustamante@puce.edu.ec

² King's College London, Department of Geography. Strand, London WC2R 2LS, United Kingdom.

³ Universidad San Francisco de Quito, College of Biological and Environmental Sciences. C.P. 17-12-841, Quito, Ecuador.

⁴ Museo Ecuatoriano de Ciencias Naturales, Sección Vertebrados, División de Herpetología Rumipamba 341 y Av. De los Shyris (Parque La Carolina), Quito, Ecuador.

⁵ Natural History Museum & Biodiversity Research Center, Department of Ecology and Evolutionary Biology, The University of Kansas. Lawrence, Kansas 66045-7561, U.S.A.

The amphibian diversity in the Chocoan tropical rain forests of Ecuador is far from being thoroughly documented. Although Neotropical equatorial forests might be the most amphibian-diverse in the world (Duellman 1978; 2005; Lynch 2005; Ron 2001–2006), few studies have been directed towards understanding the richness of the Ecuadorian Chocó.

This is especially true for Glassfrogs (Centrolenidae), which known richness in Ecuador is rapidly increasing (14 species additions to country's checklist since 2004; Guayasamin and Bonaccorso 2004; Guayasamin et al. 2006a; 2006b; Cisneros-Heredia and McDiarmid 2006; Guayasamin and Trueb

2007; Cisneros-Heredia and Meza-Ramos 2007; Cisneros-Heredia and McDiarmid 2007; Cisneros-Heredia and Yáñez-Muñoz 2007; Cisneros-Heredia 2007). Glassfrogs are mainly considered to be a montane group, and the western slopes of the Ecuadorian Andes hold the highest richness for the country with 17 described species (Cisneros-Heredia and McDiarmid 2006; Coloma 2005–2007; Cisneros-Heredia and Meza-Ramos 2007; Cisneros-Heredia and McDiarmid 2007). Some other regions, like the lowlands of the Ecuadorian Chocó, also hold high centrolenid diversity (13 species, Table 1), and its species richness could be as important as the Andean, since more species might remain undescribed.

Table 1. Centrolenid frogs occurring in Ecuadorian Chocó.

Species	Reference
<i>Centrolene callistommum</i>	Guayasamin and Trueb 2007
<i>Centrolene ilex</i>	Guayasamin et al. 2006a
<i>Centrolene litorale</i>	Grant and Morales 2004
<i>Centrolene prosoblepon</i>	Boulenger 1898
<i>Cochranella albomaculata</i>	Guayasamin et al. 2006a
<i>Cochranella mache</i>	Guayasamin and Bonaccorso 2004
<i>Cochranella ocellifera</i>	Boulenger 1899
<i>Cochranella pulverata</i>	This work
<i>Cochranella spinosa</i>	Duellman and Burrowes 1989
<i>Hyalinobatrachium aureoguttatum</i>	This work
<i>Hyalinobatrachium fleischmanni</i>	Noble 1924
<i>Hyalinobatrachium petersi</i>	Goin 1961
<i>Hyalinobatrachium valerioi</i>	Duellman and Burrowes 1989

NOTES ON GEOGRAPHIC DISTRIBUTION

Herein, we add two species to the list of Chocoan Glassfrogs in Ecuador. We report for the first time *Cochranella pulverata* and *Hyalinobatrachium aureoguttatum* from the Ecuadorian Chocó. Specimens are deposited at the Museo de Zoología, Pontificia Universidad Católica del Ecuador, Quito (QCAZ), and at the División de Herpetología, Museo Ecuatoriano de Ciencias Naturales, Quito (DHMECN).

Cochranella pulverata was known previously from lowlands to moderate elevations in the Atlantic versant of north-central Honduras and Nicaragua, Costa Rica (up to 960 m), Panama; and the Pacific versant of Costa Rica and Panama south to the Department of Nariño in Colombia (up to 300 m) (Ruiz-Carranza et al. 1996; Acosta-Galvis 2000; McCraine and Wilson 2002; Savage 2002; IUCN et al. 2006). We report the presence of *Cochranella pulverata* in the following Ecuadorian

localities (Figure 1): Provincia Esmeraldas: 1) Durango (01°03' N, 78°37' W; 100 m), on 24 May 2006, a male perching over leaf at 3 m above a creek (QCAZ 32066); 2) Río Bogotá, recinto Durango (01°00' N, 78°37' W; 120–200 m), on 18 July 2005, a vocalizing male (DHMECN 3194) and a female (DHMECN 3195), both found on epiphytes adhered to rocks about 2 m above a creek; 3) Charco Vicente (0°42' N, 78°54' W; ca. 100 m), on 15 September 1992, an adult female (QCAZ 11368); 4) Salto del Bravo, Reserva Ecológica Cotacachi Cayapas (0°40' N, 78°57' W; 100 m), on 19 April 1993, an adult female (QCAZ 11367); 5) Estero Aguacate, Parroquia San Francisco del Cabo (00°39' N, 80°03' W, 10–64 m), on 15 August 2004, a female on upper side of ferns up to 1.6 m above ground (DHMECN 2612); Province of Pichincha: 6) Silanche (00°08' N, 79°08' W; ca. 400 m), on 1 May 2006, a male calling on the upper side of a leaf, 2 m above a creek (QCAZ 32224).




Figure 1. Map showing Ecuadorian records for *Cochranella pulverata*: (1) Durango; (2) Río Bogotá; (3) Charco Vicente; (4) Salto del Bravo; (5) Estero Aguacate; and (6) Silanche.

NOTES ON GEOGRAPHIC DISTRIBUTION

Acosta-Galvis (2000) stated that the southern limit of the distribution for *Cochranella pulverata* in Colombia was Departamento Nariño (Colombian-Ecuadorian border), but did not precise a certain locality. Based on that, we extend the distributional range of the species ~ 140 km southwards, being Silanche the southern limit for the known distribution of the species. *Cochranella pulverata* was found simpatrically with *Centrolene callistommum* and *Hyalinobatrachium aureoguttatum* in Durango; with *Centrolene prosoblepon* and *H. fleischmanni* in Estero Aguacate; with *Centrolene ilex*, *C. callistommum*, *Cochranella albomaculata*, and *H. aureoguttatum*

in Río Bogotá, recinto Durango; and with *C. prosoblepon* in Charco Vicente.

Cochranella pulverata is distinguished from other Glassfrogs by the following characters: (1) lime green dorsum with small yellow-white spots; (2) light green bones; (3) venter completely transparent with white heart, liver, and digestive tract; (4) snout truncate from above and obtuse in profile; (5) indistinct tympanum; (6) SVL 22–29 mm in adult males and 23–33 mm in adult females and (7) webbing formula between outmost fingers: II 1 ½–3 III 1 ½–1 ¼ IV (Savage 2002) (Figure 2).



Figure 2. Dorsal and ventral views of *Cochranella pulverata* (QCAZ 32224).

The *Hyalinobatrachium pulveratum* species group was proposed by Ruiz-Carranza and Lynch (1991) to contain the former *Centrolenella pulverata* and *C. antisthenesi*. Cisneros-Heredia and McDiarmid (2006) proposed a new combination for both species, linking them to *Cochranella* (as *C. pulverata* and *C. antisthenesi*). We follow their proposal, with the caveat that *Cochranella*, as currently defined, is not monophyletic (Ruiz-Carranza and Lynch 1991; Guayasamin *et al.* 2006b). We suggest that further taxonomic changes in the family should be based on a phylogenetic framework to avoid subjectivity in the generic placement of species.

Hyalinobatrachium aureoguttatum was known previously from extreme southwestern Panama and the Pacific lowlands and western slopes of the Cordillera Occidental of Colombia, in the departamentos Antioquia, Chocó, Risaralda, and Valle del Cauca, at altitudes between 45–1570 m (Barrera-Rodríguez and Ruiz-Carranza 1989; Ruiz Carranza *et al.* 1996; Ibáñez *et al.* 1999; Acosta-Galvis 2000). In Ecuador, this species was collected in the following localities (Figure 3): Provincia Esmeraldas: 1) 2 km E of San Francisco (Durango - San Francisco), on the road to Durango (01°05' N, 78°41' W; 63 m), on 01 December 2004 and 25 May 2006, seven males (QCAZ 27429,

NOTES ON GEOGRAPHIC DISTRIBUTION

28802, 32101, 32105, 32129, 32132, and 32133); 2) Durango (01°03' N, 78°37' W; 100 m), on 01 July 1994 and 24 May 2006, six males (QCAZ 6302, 6303, 6441, 6442, 32069, and 32070); 3) Río Bogotá, recinto Durango (01°00' N, 78°37' W; 120–200 m), 13–15 August 2005, five males (DHMECN 3221–25); Provincia Imbabura: 4) 6 km east of Lita (00°52' N, 78°26' W; 600 m), on 01 February 1994, a male (QCAZ 4323). *Hyalinobatrachium aureoguttatum* is known to reach its southern distribution limit at 3°45' N.

Here we give the first reports of *H. aureoguttatum* for Ecuador, extending the known geographic range ~ 375 km to the south (6 km east of Lita) (Figure 3). The species was found in sympatry with *Centrolene callistommum* in 2 km east of San Francisco (Durango - San Francisco); with *Centrolene callistommum*, *Centrolene ilex*, *Cochranella albomaculata*, and *Cochranella pulverata* at Río Bogotá, recinto Durango; and with *Cochranella albomaculata* at 6 km east of Lita.




Figure 3. Map showing Ecuadorian records for *Hyalinobatrachium aureoguttatum*. (1) 2 km east of San Francisco; (2) Durango; (3) Río Bogotá; (4) 6 km east of Lita.

Hyalinobatrachium aureoguttatum is easily diagnosed by the combination of the following characters: (1) light green dorsum with a pale yellow reticulum and between two and six large, bright yellow dots (Colombian populations also have dark flecks); (2) white bones; (3) venter

completely transparent with white or red heart, and white liver and digestive tract; (4) snout rounded from above and in profile; (5) tympanic membrane fairly indistinct, tympanic annulus evident anteroventrally; and (6) SVL 19–21 mm in adult males (Figure 4).

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 4. Dorsal and ventral views of *Hyalinobatrachium aureoguttatum* (QCAZ 32068).

Acknowledgments

We are grateful to L. A. Coloma and M. Altamirano B. for granting access to specimens deposited at QCAZ and DHMECN. Pablo A. Menéndez helped with geographic information. MRB was funded by Escuela de Biología, the Pontificia Universidad Católica del Ecuador and Fundación Numashir; DFCH was supported by the Museo Ecuatoriano de Ciencias Naturales, Universidad San Francisco de Quito, M. E. and L. Heredia, the Research Training Program, and the Smithsonian Women's Committee, National Museum of Natural History (Smithsonian Institution); JMG was supported by The

University of Kansas, Fundación Numashir, and grants from the National Science Foundation (DIG DEB-0608011), and the American Philosophical Society; MYM was supported by Fundación SIRUA, Flora y Fauna Internacional and MECN; M. Herrera M. and C. Landázuri provided field support; HMOA was funded by EcoCiencia, Conservation International and the Netherlands Royal Embassy. HMOA thanks M. Larrea, manager of the Reserva Monte Saino (for providing access to facilities) and all the people involved in this effort: A. Ortiz, C. Tobar, F. Armas, M. Arellano, R. Cabrera, and F. Narváez.

Literature cited

- Acosta-Galvis, A. R. 2000. Ranas, salamandras y cecilias (Tetrapoda: Amphibia) de Colombia. *Biota Colombiana* 1(3): 289–319.
- Barrera-Rodríguez, M. and P. M. Ruiz-Carranza. 1989. Una nueva especie del género *Centrolenella* Noble 1920 (Amphibia: Anura: Centrolenidae) de la Cordillera Occidental de Colombia. *Trianea* 3: 77–84.
- Boulenger, G. A. 1898. An account of the reptiles and batrachians collected by Mr. W. F. H. Rosenberg in western Ecuador. *Proceedings of the Zoological Society of London* 1898(1): 107–126.
- Boulenger, G. A. 1899. Descriptions of new batrachians in the collection of the British Museum (Natural History). *Annals and Magazine of Natural History* 7(3): 273–277.
- Cisneros-Heredia, D. F. 2007. A new species of glassfrog of the genus *Centrolene* from the foothills of Cordillera Oriental of Ecuador. *Herpetozoa* 20 (1/2): 27–34.
- Cisneros-Heredia, D. F. and R. W. McDiarmid. 2006. A new species of the genus *Centrolene* (Amphibia: Anura: Centrolenidae) from Ecuador with comments on the taxonomy and biogeography of Glassfrogs. *Zootaxa* 1244: 1–32.
- Cisneros-Heredia, D. F. and R. W. McDiarmid. 2007. Primer registro de *Hyalinobatrachium ruedai* en

NOTES ON GEOGRAPHIC DISTRIBUTION

- Ecuador con notas sobre otras especies congenericas. *Herpetotropicos* 3(1): 21–28.
- Cisneros-Heredia, D. F. and P. Meza-Ramos. 2007. An enigmatic new species of Glassfrog (Amphibia: Anura: Centrolenidae) from the Amazonian Andean slopes of Ecuador. *Zootaxa* 1485: 33–41.
- Cisneros-Heredia, D. F. and M. H. Yáñez-Muñoz. 2007. A new species of Glassfrog (Centrolenidae) from the southern Andean foothills on the West Ecuadorian Region. *South American Journal of Herpetology* 2(1): 1–10.
- Coloma, L. A. (ed.). 2005–2007. Anfíbios de Ecuador. [en línea]. Ver. 2.0 (29 Octubre 2005). Museo de Zoología, Pontificia Universidad Católica del Ecuador. Quito, Ecuador. Electronic Database accessible at <http://www.puce.edu.ec/zoologia/vertebrados/amphibiawebec/index.html>. Captured on August 2007.
- Duellman, W. E. 1978. The biology of an Equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous publications of the University of Kansas, Museum of Natural History* 65: 1–352.
- Duellman, W. E. 2005. Cusco Amazónico, the lives of amphibian and reptiles in an Amazonian rainforest. Ithaca, Cornell University Press. 433 p.
- Duellman, W. E. and P. A. Burrowes. 1989. New species of frogs, *Centrolenella*, from the pacific versant of Ecuador and southern Colombia. *Occasional Papers of the Museum of Natural History, University of Kansas* 132: 1–14.
- Goin, C. J. 1961. Three new centrolenid frogs from Ecuador. *Zoologischer Anzeiger* 166: 95–104.
- Grant, T. and M. Morales. 2004. *Centrolene litorale*. 2006 IUCN Red List of Threatened Species. Electronic Database accessible at <http://twwww.iucnredlist.org>. Captured on March 2007.
- Guayasamin, J. M. and E. Bonaccorso. 2004. A new species of glass frog (Centrolenidae: *Cochranella*) from the lowlands of northwestern Ecuador, with comments on the *Cochranella granulosa* group. *Herpetologica* 60(4): 485–494.
- Guayasamin, J. M. and L. Trueb. 2007. A new species of Glass Frog (Anura: Centrolenidae) from the lowlands of northwestern Ecuador, with comments on centrolenid osteology. *Zootaxa* 1447: 27–45.
- Guayasamin, J. M., D. F. Cisneros-Heredia, M. Yáñez-Muñoz, and M. Bustamante. 2006a. Notes on geographic distribution. Amphibia, Centrolenidae, *Centrolene ilex*, *Centrolene litorale*, *Centrolene medemi*, *Cochranella albomaculata*, *Cochranella ametarsia*: Range extensions and new country records. *Check List* 2(1): 24–25.
- Guayasamin, J. M., M. R. Bustamante, D. Almeida-Reinoso, and W. C. Funk. 2006b. Glass frogs (Centrolenidae) of Yanayacu Biological Station, Ecuador, with the description of a new species and comments on centrolenid systematics. *Zoological Journal of the Linnean Society* 147: 489–513.
- Ibáñez, R., F. Jaramillo, and C. Jaramillo. 1999. Ampliación del ámbito de distribución y descripción del renacuajo de la rana de cristal *Hyalinobatrachium aureoguttatum* (Anura: Centrolenidae). *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 23: 293–298.
- IUCN, Conservation International, and NatureServe. 2006. Global Amphibian Assessment. Electronic Database accessible at <http://www.globalamphibians.org>. Captured on August 2007.
- Lynch, J. D. 2005. Discovery of the richest frog fauna in the world — An exploration of the forests to the north of Leticia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 113: 581–588.
- McCrane, J. R. and L. D. Wilson. 2002. The Amphibians of Honduras. Ithaca, Society for the Study of Amphibians and Reptiles. 623 p.
- Noble, G. K. 1924. Some Neotropical batrachians preserved in the United States National Museum with a note on secondary sexual characters of these and other amphibians. *Proceedings of the Biological Society of Washington* 37: 65–72.
- Ron, S. R. 2001–2006. Anfíbios de Parque Nacional Yasuní, Amazonía ecuatoriana. [online]. Ver. 1.4 (2006). Museo de Zoología, Pontificia Universidad Católica del Ecuador. Quito, Ecuador. Electronic Database accessible at <http://www.bio.utexas.edu/grad/ecuador/web/yasuni/esp/anfyas.htm>. Captured on August 2007.
- Ruiz-Carranza, P. M. and J. D. Lynch. 1991. Ranas Centrolenidae de Colombia I: propuesta de una nueva clasificación genérica. *Lozania* 57: 1–30.
- Ruiz-Carranza, P. M., M. C. Ardila-Robayo, and J. D. Lynch. 1996. Lista actualizada de la fauna de amphibia de Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 20(77): 365–415.
- Savage, J. M. 2002. The Amphibians and Reptiles of Costa Rica: An Herpetofauna between two Continents, between two Seas. Chicago, University of Chicago Press. 934 p.

Received April 2007

Accepted September 2007

Published online September 2007